



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE ON MULTIPLY PERFECT NUMBERS

BY R. D. CARMICHAEL

THE object of this note is to prove the following proposition :

There exist no multiply perfect numbers of multiplicity 3 of the form $m = p_1^{a_1} p_2^{a_2} p_3^{a_3}$ where p_1, p_2, p_3 are distinct primes and $p_1 < p_2 < p_3$.*

In the ANNALS OF MATHEMATICS, ser. 2, vol. 2, p. 172 (July 1901), Dr. Jacob Westlund shows that $p_1 = 2$ and $p_2 = 3$. Hence we have (cf. Westlund, *l. c.*), supposing that there are such multiply perfect numbers,

$$(1) \quad 3 = \frac{2^{a_1+1} - 1}{2^{a_1}} \cdot \frac{3^{a_2+1} - 1}{3^{a_2} \cdot 2} \cdot \frac{p_3^{2a_3+1} - 1}{p_3^{2a_3}(p_3 - 1)},$$

from which we get

$$(2) \quad p_3^{2a_3} + p_3^{2a_3-1} + \dots + p_3 + 1 = \frac{p_3^{2a_3} \cdot 2^{a_1+1} \cdot 3^{a_2+1}}{(2^{a_1+1} - 1)(3^{a_2+1} - 1)}.$$

The first member is not divisible by p_3 ; hence,

$$(3) \quad p_3^{2a_3} + p_3^{2a_3-1} + \dots + p_3 + 1 = 2^\lambda 3^\mu,$$

where λ and μ are to be determined and belong to the series $0, 1, 2, 3, \dots$.

Since p_3 is a prime greater than 3 it is of one of the forms $6n \pm 1$.

Case I. Let $p_3 = 6n - 1$. Substituting in equation (3), we have

$$(4) \quad (6n - 1)^{2a_3} + (6n - 1)^{2a_3-1} + \dots + 1 = 2^\lambda 3^\mu.$$

If each term of the first member is expanded, the sum of the resulting terms not containing $6n$ as a factor is 1. Hence, this first number is divisible by neither 2 nor 3. This case therefore cannot yield multiply perfect numbers.

* This term was introduced by D. N. Lehmer in 1901; see ANNALS OF MATHEMATICS, ser. 2, vol. 2, p. 103: "a multiply perfect number is one which is an exact divisor of the sum of all the divisors, the quotient being the multiplicity."

Case II. Let $p_3 = 6n + 1$. Substituting in equation (3) after summing its left member, we have

$$(5) \quad \frac{(6n+1)^{2a_3+1} - 1}{6n} = 2^\lambda 3^\mu,$$

whence

$$(6) \quad (6n)^{2a_3} + (2a_3 + 1)(6n)^{2a_3-1} + \dots + a_3(2a_3 + 1)(6n) + (2a_3 + 1) = 2^\lambda 3^\mu.$$

Let $2a_3 + 1 = 3^\nu \cdot t$, where t contains neither 2 nor 3 as a factor. Each term but the last of the first member of (6) is divisible by $2 \cdot 3^{\nu-1}$. Hence, dividing through by 3^ν , equation (6) may be written (q being an integer quotient),

$$(8) \quad 6 \cdot q + t = 2^\lambda 3^{\mu-\nu}.$$

The first member of (8) is greater than 1 and is divisible by neither 2 nor 3. Hence, equation (8) being impossible, case II yields no multiply perfect numbers.

Hence the proposition.